# REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this

And the second	2. REPORT DATE	3. REPORT TYPE AND I	TYPE AND DATES COVERED	
1. AGENCY USE ONLY (Leave blank)	1 Jan 98	Final 1 Ja	nuary 90 to 31 Dec. 95	
4. TITLE AND SUBTITLE  Exposure of human C fi	cells to electrom	agnetie	5. FUNDING NUMBERS NO0074.90 . J - 1266	
6. AUTHORISI Ann 5. Henderson		(190 <u>0)</u> 10 (10 (10)	\$ 100 miles   100	
7. PERFORMING ORGANIZATION NAME(	S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION REPORT NUMBER	
Hunter College - CUA 695 Park Avenue New York ny	10021 1003.	And the second of the second o	A)A  10. SPONSORING / MONITORING	
9. SPONSORING / MONITORING AGENC Office of Naval 800 N QUINCY	Research, Code	WI SB	AGENCY REPORT NUMBER	
Arlington UA 27	2217		1:	
11. SUPPLEMENTARY NOTES				
120. DISTRIBUTION / AVAILABILITY ST.	•	er en	12b. DISTRIBUTION CODE	

#### 13. ABSTRACT (Maximum 200 words)

The objective of this research was to find how exposure to extremely low frequency (elf) electric and magnetic fields (EMFs) affects human cells. It was previously postulated that EMFs initiate a transduction cascade at the cell membrane. If this is the case, then within minutes, a sequence of events, including early gene activation, could set up the initial steps leading to proliferation. A series of experiments were set up to determine the validity of this model by measuring specific proteins or protein groups. The results showed that some proteins are increased following exposure to a 60 Hz EMF for periods from 45 to 120 minutes. Other proteins, however, are either decreased or unaffected by the presence of the field.

14. SUBJECT TERMS		15. NUMBER OF PAGES
Electromagnetia Fields		16. PRICE CODE
	CLASSIFICATION 19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
OF REPORT OF THIS P.	Ade (	Standard Form 298 (Rev. 2-89) Standard Form 298 (Rev. 2-89)
	•	Standard Form 290 (NS-Std Z39-18 Prescribed by ANSI-Std Z39-102

### FINAL REPORT

Grant #: N00014-90-J-1266

PRINCIPAL INVESTIGATOR: Ann S. Henderson

INSTITUTION: Hunter College-CUNY

GRANT TITLE: Exposure of Human Cells to Electromagnetic

Fields

AWARD PERIOD: 1 January 1990 - 31 December 1995

OBJECTIVES: To find how exposure to extremely low frequency (elf) electric and magnetic fields (EMFs) effects human cells. It is postulated that EMFs initiate a signal transduction cascade at the cell membrane leading to changes in the influx of calcium and/or receptor binding. If this is the case, then within minutes, a sequence of events, including early gene activation, could set up the initial steps leading to proliferation. A series of experiments has been set up to determine the validity of this model. The aims are to make specific comparisons between unexposed and exposed cells which include measuring protein synthetic patterns. The hypothesis that common pathways are used by cells exposed to EMFs or TPA is also tested in this study. Inherent in our approach is to develop unequivocal methods for measurement of the effects of EMFs.

<u>APPROACH</u>: HeLa or HL-60 cells are exposed to 60 Hz sinusoidal EMFs at 8  $\mu T$  for 20 minutes. Different experimental approaches have been used to investigate the effect of EMF exposure on polypeptide synthetic activity, enzymatic activity or protein modification. The experimental designs include setting conditions by measurement of products of transfected genes, immunoprecipitation to measure protein phosphorylation, inhibition of enzymatic activity and determinations using 2-D gel electrophoresis.

ACCOMPLISHMENTS. (1) Phosphorylation of Fos protein. The Fos protein becomes phosphorylated at serine residues present at the carboxyl terminus of the protein. Other studies have shown that the phosphorylation of the protein, causes the protein [in conjugation with Jun] to down-regulate its own promoter. EMF (or TPA) exposure results in an increase in the amount of modified (phosphorylated) Fos protein, with a maximum increase of 50 % following 20 minutes exposure. (2) Expression of CAT protein in transfected HeLa cells. Another approach is to measure CAT (linked to an appropriate upstream regulatory region of the

c-fos or c-myc gene) protein following transfections of HeLa cells. CAT protein levels are then compared between unexposed and exposed cells; TPA exposure is used alone, as well in combination with EMF. CAT protein is maximally expressed at one hour following exposure, and this is also the time at which the maximum difference between CAT protein in unexposed and exposed cells is achieved. Irrespective of the order of exposure of cells to TPA or EMF in dual exposures, the effects were not synergistic. (3) The role of PKC. The most compelling evidence for the validity of a given observation is that the effect can be erased by inhibition. PKC was inhibited by chelerythreine (1  $\mu M$ ). In the presence of the inhibitor, there was no increase in CAT protein in transfected HeLa cells exposed to EMFs. The ratio of membrane to cytoplasmic PKC $\alpha$  levels was compared in HL-60 cells with no exposure or exposed to TPA or EMFs. There is an increase observed at 10 minutes following either TPA treatment or EMF exposure. Although the effect of EMF exposure is much smaller that using TPA, the kinetics are the same. (4) Optimizing conditions for studying proteins in cells exposed to EMFs. Analysis of 2-D gels is by identification of known polypeptides and "grid analysis" of total gels using computer assistance. The results showed increases or decreases in some polypeptides from exposed cells. There was, however, variability in the results (based on control studies) which prompted us to analyze our system more carefully. Our previous protocol used a time period of 25 minutes following exposures. This turned out to be inadequate, based on studies of specific genes. We then did a very precise study over time as to when differences in specific proteins (Fos and Myc) were maximized following exposure to EMFs. The results showed that a period of one hour (following exposure) was required.

CONCLUSIONS: These studies: (a) developed appropriate conditions for determination of changes in specific proteins that are quantitatively changed in the presence of EMFs and (b) suggest that EMFs can effect the regulation of c-fos by post- translational modification of the gene product.

SIGNIFICANCE: The studies show that elf EMF exposure can affect proteins required in cellular regulatory pathways.

PATENT INFORMATION: NONE

AWARD INFORMATION: NONE

# PUBLICATIONS AND ABSTRACTS (total period of grant):

### Publications

- 1. Goodman, R. and A. Henderson, 1991. Transcription in cells exposed to extremely low frequency electromagnetic fields: a review. Bioelectrochem. Bioenerget 25, 335-355.
- 2. Goodman, R., J. Bumann, L-X Wei, J-C Xu and A. Shirley-Henderson, 1991. Transcriptional changes in cells exposed to extremely low frequency electromagnetic fields. In Electromagnetics in Biology and Medicine (Carl T. Brighton and Soloman R. Pollack, eds), pp. 127-132, San Francisco Press, San Francisco.
- 3. Hubbard-Smith, K., P. Patsalis, J. Pardinas, J. Jha, A. Henderson and H. Ozer 1992. Altered chromosome 6 in immortal human fibroblasts. Mol. Cell Biol., 12, 2273-2281.
- 4. Goodman, R., D. Weisbrot, A. Uluc and A. Henderson, 1992. Transcription in Drosophila melanogaster salivary gland cells is altered following exposure to low frequency electromagnetic fields: Analysis of chromosome 3R. Bioelectromagnetics 13: 111-118.
- 5. Blank M., L. Soo, H. Lin, A. Henderson and R. Goodman, 1992. Changes in transcription in HL-60 cells following exposure to alternating current from electric fields. Bioelectrochem. Bioenerget. 28:301-309.
- 6. Goodman, R., D. Weisbrot, A. Uluc and A. Henderson, 1992 Transcription in Drosophila melanogaster salivary gland cells is altered following exposure to low frequency electromagnetic fields: Analysis of chromosomes 3L and X. Bioelectrochem. Bioenerget., 28: 311-318.
- 7. Goodman, R., J. Bumann, L-X Wei and A. Shirley-Henderson, 1992. Exposure of human cells to electromagnetic fields: Effect of time and field strength on transcript levels.
- 8. Goodman, R., L-X Wei, J. Bumann and Ann S. Henderson, 1992. Exposures to electric and magnetic (EM) fields increases transcripts in HL-60 cells; does adaptation to EM fields occur? Bioelectrochem Bioenergetics 29: 185-192.
- 9. Weisbrot, D., Uluc, A., Henderson, A. and R. Goodman, 1993. Transcription in Drosophila melanogaster salivary gland cells is altered following exposure to low frequency electric and magnetic fields: analysis of chromosomes 2R and 2L. Bioelectrochem. Bioenerget. 31:39-47.
- 10. Goodman, R., Yuri Chizmadzhev and Ann Shirley-Henderson, 1993. Electromagnetic fields and cells. J. Cellular Biochemistry 52: 1-6.
- 11. Weisbrot, D.R., O. Khorkova, H. Lin, A. Henderson and R. Goodman, 1993. The effect of low frequency electric and magnetic fields on gene expression in Saccharomyces cerevisiae. Bioelectrochem. Bioenerget. 31: 167-177. 12. Blank, M., L. Soo, H. Lin, R. Goodman and A.S. Henderson,

- 1993. Stimulation of transcription in HL-60 cells by alternating currents from electrical fields. In Electricity and Magnetism in Biology and Medicine (M. Blank, ed.) San Francisco Press, San Francis, pp. 516-517.
- 13. Broude, N., R. Karabakhtsian, N. Shalts, R. Goodman and A. Henderson, 1994. Correlation between the amplitude of plasma fluctuations and the response of cells to electric and magnetic (EM) fields. Bioelect. Bioenerget. 31: 19-23. 14. Lin, H., R. Goodman and A. Henderson, 1994. Specific
- 14. Lin, H., R. Goodman and A. Henderson, 1994. Specific region of the c-myc promoter is responsive to electric and magnetic fields. J. Cell. Biochemistry 55:1-8.
- 15. Goodman, R., M. Blank, H. Lin, O. Khorkova, L. Soo, D. Weisbrot and A. Henderson, 1994. Increased levels of hsp70 transcripts induced when cells are exposed to low frequency electromagnetic fields. Bioelectrochem. Bioenerget.33: 115-120.
- 16.Hadjiargyrou, M., K. McDowell and A. Henderson, 1994. A transfected human ribosomal RNA gene is active in nucleolus formation in human cells. Cytogenet. Cell Genet. 60: 58-62. 17. McDowell, K., M. Hadjiargyrou, P. Patsalis and A. Henderson, 1994. A partial human rDNA repeat functions as a nucleolar organizer region. Cytogenet. Cell Genet. 60: 63-67.
- 18. Gold, S., R. Goodman and A. Henderson, 1994. Exposure of human cells to electric and magnetic fields results in increased levels of T-antigen mRNA and protein. Bioelectromagnetics 15: 329-336.
- 19. Goodman, R. and A. Henderson, 1994. Effects of electric and magnetic fields on gene expression. In Biological effects of electromagnetic fields (D. Carpenter, ed.) Academic Press, Orlando, FLA, pp.155-176.
- 20. Karabakhtsian, R., N. Broude, N. Shalts, S. Kochlatyi, R. Goodman and A. S. Henderson, 1994. Calcium is necessary in the cell response to EM fields. FEBS LETT. 349: 1-6.
  21. Patsalis, P.C., H.L. Ozer and A. Henderson, 1995.
  Sequence and STS of two new single copy DNA markers p21.6A
- (D6S116) and p21.14A (D6S122) on chromosome region 6q14-15; D6S122 highly homologous to the IgK chain. Clinical Genetics 48:275-276.
- 22. Rao, S. and A. Henderson, 1996. Regulation of c-fos is affected by electromagnetic fields. J. Cell Biochem. 63: 358-365.

#### Abstracts and Presentations

1. Goodman, R., L-X Wei and A. Henderson, 1991. Relationship of fields strength and time to induction of increased transcripts by elf EM fields. RCMI Symposium, Atlanta.
2. Goodman, R. and A. Henderson, 1991. Transcriptional changes in cells exposed to extremely low frequency

- electromagnetic fields. FASEB, Atlanta,
- 3. Goodman, R., J. Bumann and A. Henderson (1991) Defining signal parameters which induce transcriptional/translational alterations in HL-60 cells. abstract presented at 13th Ann Bioelectromagnetics Soc. Salt Lake City, B-2-3.
- 4. Patsalis, P., K. Hubbard-Smith, J. Pardinas, K. Jha, A. Henderson and H. Ozer, 1991. Deletion of 6q is involved in immortalization of SV-40 transformed human fibroblasts. 8th International Congress of Human Genetics, Washington.
- 5. Goodman, R., L-X Wei, S. Gold and A. Henderson, 1991. Transcriptional and translational alterations in human cells are induced by low frequency electromagnetic fields: Relationship to time and exposure parameters. American Society of Cell Biology, Boston, MA.
- 6. Goodman, R., L-X Wei and A. Henderson, 1991. An increase in mRNA and protein levels is induced in human cells exposed to electromagnetic fields: relationship to time of exposure, post-exposure time and frequency. DOE/Contractor's Review. Milwaukee, WI, A-18
- 7. Goodman, R., L-X Wei, J. Bumann and A. Henderson, 1991. Increased transcripts in HL-60 cells exposed to EM fields: Comparison of exposure geometries. DOE/Contractor's Review. Milwaukee, WI, p-13
- 8. Goodman, R., L-X Wei, S. Gold and A. Henderson, 1992. An increase in messenger RNA and protein levels is induced in cells exposed to electromagnetic fields: the relationship of time and exposure conditions. World Congress, Bioelectromagnetics, Orlando.
- 9. Goodman, R., J. Bumann, L-X Wei and A. Henderson, 1991. Increased transcripts in HL-60 cells exposed to EM fields: comparison of exposure geometries. The Ann. Rev. of Research on Biological Effects on 50 and 60 Hz Electric and Magnetic Fields, Milwaukee, WI, P-13.
- 10. Camargo, E., S. Ogiste, R. Goodman and A. Henderson, 1991. Effect of 60 Hz EM fields on cell division in human HL-60 cells. RCMI Symp, Washington, 1991.
- 11. Bumann, J., R. Goodman and A. Henderson, 1991. Die Wirkung Niederfrequenter Elektromagnetischer Wellen auf die Gentranskription, Stuttgart
- 12. Blank, M., L. Soo, H. Lin, A. Henderson and R. Goodman, 1991. Stimulation of transcription in HL-60 cells by alternating currents from electric fields. DOE/Contractors Review, Milwaukee, WI, A-19
- 13. Goodman, R., L-X Wei and A. Henderson, 1992. An increase in mRNA and protein levels is induced in human cells exposed to electromagnetic fields: Relationship to time and exposure conditions. First World Congress for Electricity and Magnetism in Biology and Medicine, Orlando, FL.
- 14. Goodman, R., L-X Wei and A. Henderson, 1992. Changes in translational patterns are induced in human HL-60 cells at

- specific times and field strengths. First World Congress for Electricity and Magnetism in Biology and Medicine, Orlando, FL.
- 15. Goodman, R, O. Khorkova, A. Henderson and D.Weisbrot, 1992. The effect of low frequency electric and magnetic fields on gene expression in Saccharomyces cerevisiae. DOE Contractor's Meeting, San Diego, p.15.
- 16. Goodman, R. M. Blank, O. Khorkova, D. Weisbrot, and A. Henderson, 1992. The heat shock hsp70 gene is induced in a variety of cell types exposed to elf EM fields or ac current. Does the cell respond to EM field exposure as a cellular stress? DOE Contractor's Meeting, San Diego, A1 17. Gold, S., R. Goodman and A. Henderson, 1992. Increased levels of T-antigen mRNA and protein in human cells exposed to 60 Hz sinusoidal field. DOE Contractor's Meeting, San Diego.
- 18. Lin, H., R. Goodman and A. Henderson. EM field exposure affects the c-myc promotor. DOE Contractor's Meeting, San Diego.
- 19. Patsalis, P., H. Ozer and A. Henderson, 1992. A breakpoint on chromosome 6q is involved in cell immortalization. Chromosomes in Cancer, Phoenix, Az.
- 20. Lin, H., R. Goodman and A. Shirley-Henderson, 1992. The role of transcriptional regulation in the response of cells to EM fields. DOE Contractor's Meeting, San Diego.
- 21. Weisbrot, D., O. Khorkova, A. Henderson and R. Goodman, 1993. Transcript levels for some genes, including a heat shock gene, are increased in Saccharomyces cerevisiae cells exposed to elf EM fields. BEMS, Los Angeles, p.40.
- 22. Goodman, R., M. Blank, O. Khorkova, D. Weisbrot, and A. Henderson, 1993. Increased expression of heat shock gene hsp70 is measured in cells exposed to electric and magnetic fields: is em field exposure a cellular stress? BEMS, Los Angeles, p.4.
- 23. Lin, H., R. Goodman and A. Henderson., 1993. EM field exposure affects the c-myc promotor. BEMS, Los Angeles, p.41.
- 24. Lin, H., R. Goodman and A. Henderson, 1993. DNA upstream of the c-myc gene is responsive to electric and magnetic fields. Annual Review of Research on Biological Effects of Electric and Magnetic Fields from the Generation, Delivery and Use of Electricity, Savannah, Ga., pp.11-12
- 25. Leacock, J., S. Rao and A. Henderson, 1993. Effects of electromagnetic fields on HL60 cells. Presented at National MBRS symposium in Atlanta (November, 1993)
- 26. Karabakhtsian, R., N. Broude, N. Shalts, R. Goodman and A. Henderson, 1994. The presence of calcium is important in the cell's response to EM fields. BEMS, Copenhagen.
- 27. Rao, S., H. Lin, R. Goodman and A. Henderson, 1994. Regulatory DNA upstream of the c-fos and c-myc genes is

- responsive to electric and magnetic fields. BEMS, Copenhagen.
- 28. Goodman, R., M. Blank, H. Lin, L. Soo, D. Weisbrot and A. Henderson, 1994. Increased levels of hsp70 transcripts are induced when cells are exposed to low frequency electromagnetic fields. BEMS, Copenhagen
- 29. Goodman, R. H. Lin and A. Henderson, 1994. Increased transcripts for hsp70 and c-myc in cells re-stimulated with different exposure conditions. BEMS, Copenhagen
- 30. Patsalis, P., H. Ozer and A. Henderson, 1994. Primary and secondary abnormalities associated with cellular senescence and immortalization in HDFs transformed by SV40. Hellenic Association of Medical Geneticists, Thessaloniki, Greece.
- 31. Kochlatyi, S., N. Shalts, R. Karabakhtsian and A. S. Henderson, 1994. The role of transmembrane calcium flux on EMF effects on cells. DOE Contractor's Meeting, Albuquerque.p. 82
- 32. Goodman, R., H. Lin, S. Rao, R. Karabakhtsian, N. Shalts, S. Kochlatyi and A. S. Henderson, 1994. Tests of EMF results; effects can be blocked. DOE Contractor's Meeting, Albuquerque. P. 10.
- 33. Goodman, R and A. S. Henderson, 1994. What constitutes replication? DOE Contractor's Meeting, Albuquerque. P. 68 34. Rao, S. and A. Henderson, 1995 Regulatory DNA upstream of the c-fos gene is responsive to EMF exposure, BEMS, Boston. P.2
- 35. Rao, S. and A. Henderson, 1995. The Sre/Ap-1 region of the c-fos promoter is responsive to EMF; correlation of the effects of TPA and EMF. DOE Contractor's Meeting, Palm Springs. P. 11
- 36. Rao, S. and A. Henderson, 1996. Common cellular pathways used by TPA and EMF exposure? BEMS, Victoria 37. Kochlatyi, S., S. Rao, N. Shalts and A. Henderson, 1996. Designing repetition experiments; search for internal

controls. BEMS, Victoria